



See Package Insert for further directions on use of the BD® 10 mL syringe with TwinPak™ Dual Cannula Device.



1. WITHDRAW 10 mL of diluent using the BD® 10 mL syringe with TwinPak™ Dual Cannula Device included in the kit. Use only the diluent provided in this kit.



2. INJECT entire contents into the TNKase® (Tenecteplase) vial, directing the diluent at the powder. Slight foaming is common.




3. GENTLY SWIRL vial contents are completely dissolved. **DO NOT SHAKE.** Reconstitution should be complete in approximately 1 minute. Solution should be colorless or pale yellow and transparent. **USE UPON RECONSTITUTION.** If not used immediately, refrigerate solution at 2-4°C and use within 8 hours.



4. INSPECT the solution visually for particulate matter or discoloration. **WITHDRAW** the appropriate volume of solution based on patient weight. (See Dosing Information.) Discard solution remaining in the vial.

5. PRECIPITATION may occur when TNKase® (Tenecteplase) is administered in an IV line containing dextrose. To prevent precipitation, flush a dextrose-containing line with a saline-containing solution prior to and following single-bolus administration of TNKase.

ADMINISTER as an intravenous BOLUS over 1 second.



Tenecteplase Kit with:

- Neon tenecteplase sticker OVER the MI dosing table
- 5mL syringe in each kit
- Tenecteplase dosing card